

On connection between division sequences and presentations of a free group

Natalia Vanetik

CS department, Ben-Gurion University, Beer-Sheva, Israel
orlovn@cs.bgu.ac.il

Abstract. This paper describes a simple method for estimating lower bounds on the number of classes of equivalence for a special kind of integer sequences, called division sequences. The method is based on adding group structure to classes of equivalence and studying properties of resulting groups as presentations of free group.

1 Division sequences

Let $M \in \{\mathbb{Z}_{\{\neq 0\}}, \mathbb{Z}_{\{> 0\}}\}$ be a domain of mapping $\mathcal{C}_{p,q}^M : M \rightarrow M$ defined as follows.

$$\mathcal{C}_{p,q}^M(c) = \begin{cases} pc + 1, & q \text{ does not divide } c \\ c/q, & \text{otherwise} \end{cases} \quad (1)$$

We call the mapping $\mathcal{C}_{p,q}^M$ *division sequence* for parameters p and q and domain M .

Division sequence $\mathcal{C}_{p,q}^M$ induces following equivalence relation on M : for $c, d \in M$ we say that c is *equivalent* to d , denoted $c \sim d$, if $d = \mathcal{C}_{p,q}^M(c)$; symmetry, reflexivity and transitivity are added to $\sim$ in order for it to be a proper equivalence relation.

2 Presentations of free group

Let us observe monoid $M \in \{\mathbb{Z}_{\{\neq 0, \cdot\}}, \mathbb{Z}_{\{> 0, \cdot\}}\}$ and corresponding Grothendieck's group $F := K(M) \in \{\mathbb{Q}_{\{\neq 0, \cdot\}}, \mathbb{Q}_{\{> 0, \cdot\}}\}$.

We are interested in free group F with countably many generators (primes) and defining relations that (a) include abelinizing relation and (b) reflect equivalence relation $\sim$ defined on M by $\mathcal{C}_{p,q}^M$.

For fixed $p, q \in M$ we observe following relations:

$$q = 1, \quad (2)$$

$$P \sim Q \Rightarrow P = Q \quad (3)$$

For given p and q , we denote relations generated by (3) by $R_{p,q}$. Abelinizing relation that we add to F is denoted $XY = YX$ and it implies that $xy = yx$ for all pairs of generators x, y of F . Note that $R_{p,q}$ is a countable set.

Let

$$H_{p,q} := \langle F \mid XY = YX, q = 1, R_{p,q} \rangle$$

be presentation of F (we write F instead of F 's generators for simplicity). We denote by $\phi_{p,q}$ the corresponding free group homomorphism defining $H_{p,q}$. If the choice of F needs to be clarified for $H_{p,q}$, we write $H_{p,q}^M$ for domain M and free group $F = K(M)$.

Property 1. $H_{p,q}$ is isomorphic to a quotient group of H_{p,q^n} for all $n \in \mathbb{N}$.

Proof. By Tietze transformations rules (see, e.g., [Lyndon, Schupp 2001]), implication $q = 1 \Rightarrow q^n = 1$ ensures that $H_{p,q}$ is isomorphic to a quotient group of H_{p,q^n} . Since natural homomorphism $\chi : H_{p,q} \rightarrow H_{p,q^n}$, where $\chi(a) = a$, is surjective we have $\text{Im } \chi \trianglelefteq H_{p,q^n}$. $\square$

When we are interested in parameters p, q for which $H_{p,q} \cong \{1\}$, i.e. is a trivial group, following properties are obvious.

$$\begin{aligned} H_{p,q} \cong \{1\} &\implies H_{p,q^n} \cong \{1\} \text{ and} \\ H_{p,q} \not\cong \{1\} &\implies H_{p,q^n} \not\cong \{1\} \forall n \in \mathbb{N} \end{aligned} \quad (4)$$

Another indication of structure of $H_{p,q}$ are groups $H_{pp',q}$, where $p' \in F$.

Property 2. Let $p' \in \text{Ker } \phi_{p,q}$. Then $H_{p,q}$ is isomorphic to a quotient group of $H_{pp',q}$.

Proof. Let $pp'P + 1 = Q$ imply defining relation $P = Q \in R_{pp',q}$. Since $H_{p,q}$ and $H_{pp',q}$ share domain, $pp'P + 1 = Q$ also implies that defining relation $p'P = Q \in R_{p,q}$. Relations $p'P = Q, p' = 1 \in R_{p,q}$ give us defining relation $P = Q \in R_{p,q}$ by Tietze transformations rules. Therefore, each defining relation of $H_{pp',q}$ is also a defining relation of $H_{p,q}$, and we have the property. $\square$

Tietze transformations allow us to make a stronger claim under following restrictions.

Property 3. Let $q' \in \text{Ker } \phi_{p,q} \cap \text{Ker } \phi_{p,qq'}$. Then

- (a) $H_{p,q}$ is isomorphic to a quotient group of $H_{p,qq'}$,
- (b) $H_{p,qq'}$ is isomorphic to a quotient group of $H_{p,q}$.

Proof. Condition $q' \in \text{Ker } \phi_{p,q} \cap \text{Ker } \phi_{p,qq'}$ ensures that

$$q = 1, q' = 1, qq' = 1 \in R_{p,q} \cap R_{p,qq'} \quad (5)$$

Let us assume that equality

$$pP + 1 = q^n (q')^m Q \quad (6)$$

holds in $\mathbb{Q}$. Then $pP + 1 = (q')^m Q$ holds in $\mathbb{Q}/\langle q=1 \rangle$ and thus

$$P = (q')^m Q \in R_{p,q} \Rightarrow P = Q \in R_{p,q} \quad (7)$$

On other hand, equality (6) gives us defining relations

$$P = q^{n-\min(m,n)} (q')^{m-\min(m,n)} Q \in R_{p,qq'} \Rightarrow P = Q \in R_{p,qq'} \quad (8)$$

by Tietze transformations rules because $q' = 1, q = 1 \in R_{p,qq'}$. Therefore, $H_{p,q}$ is isomorphic to a quotient group of $H_{p,qq'}$ and vice versa. $\square$

Corollary 1. *If one of $H_{p,q}$, $H_{p,qq'}$ is finite and Property 3 holds for q' , $H_{p,q} \cong H_{p,qq'}$ by cardinality argument. In particular, $H_{p,q} \cong \{1\}$ if and only if $H_{p,qq'} \cong \{1\}$. $\square$*

Since moving from group $\mathbb{Q}_{\{\neq 0, \cdot\}}$ to group $\mathbb{Q}_{\{> 0, \cdot\}}$ in terms of defining relations can be done by adding defining relation $-1 = 1$, we also have

$$H_{p,q}^{\mathbb{Q}_{\{> 0, \cdot\}}} \text{ is isomorphic to a quotient group of } H_{p,q}^{\mathbb{Q}_{\{\neq 0, \cdot\}}} \quad (9)$$

Let us now observe group $\text{Ker } \phi_p$ which is a subgroups of F . Since $\text{Ker } \phi_{p,q} \subseteq \text{Ker } \phi_{p,q^n}$ by Property 1, we have normal subgroup relation

$$\text{Ker } \phi_{p,q} \trianglelefteq \text{Ker } \phi_{p,q^n} \quad (10)$$

Then following corollaries are implied by Properties 2-3.

Corollary 2. *Let $p' \in \text{Ker } \phi_{p,q}$. Then $\text{Ker } \phi_{pp',q} \trianglelefteq \text{Ker } \phi_{p,q}$. $\square$*

Corollary 3. *Let $q' \in M$ such that $q' \in \text{Ker } \phi_{p,q} \cap \text{Ker } \phi_{p,qq'}$. Then $\text{Ker } \phi_{p,q} = \text{Ker } \phi_{p,qq'}$ and $\text{Ker } \phi_{p,q} \cong \text{Ker } \phi_{p,qq'}$. $\square$*

Structure of $\text{Ker } \phi_{p,q}$, if nontrivial, can be partially disclosed as follows. Let us fix $p, q \in M$ and define

$$\overline{R}_{p,q} := P \approx 1 \Rightarrow P = 1 \quad (11)$$

to be set of defining relations, where $\sim$ is equivalence relation of a division sequence $\mathcal{C}_{p,q}^M$.

Now we can define group $\overline{H}_{p,q}$ as representation of free group F as follows.

$$\overline{H}_{p,q} := \langle F \mid XY = YX, q = 1, \overline{R}_{p,q} \rangle \quad (12)$$

We denote the corresponding homomorphism of F by $\overline{\phi}_{p,q}$. Since $\text{Im } \overline{\phi}_{p,q} \subseteq \text{Ker } \phi_{p,q}$, the following holds.

Property 4. $\overline{H}_{p,q}$ is isomorphic to a quotient group of $\text{Ker } \phi_{p,q}$. $\square$

Property 5. Let $p', q' \in M$. Then $\overline{H}_{p,q}/\langle p'=1 \rangle$ is isomorphic to a quotient group of $\overline{H}_{pp',q}$ and $\overline{H}_{p,q}/\langle q'=1 \rangle$ is isomorphic to a quotient group of $\overline{H}_{p,qq'}$.

Proof. Equivalence relation $P \sim Q$ implied by equality $pp'P + 1 = Q$ generates defining relation $P = Q \in R_{pp',q}$, defining relation $p'P = Q \in R_{p,q}$ in $H_{p,q}$ and defining relation $P = Q$ of $H_{p,q}/\langle p'=1 \rangle$. Therefore relations $P \sim Q \in \overline{R}_{pp',q}$, where $P, Q \approx 1$, generate defining relation $P = Q \in R_{p,q}$ and therefore are defining relations of $\overline{H}_{p,q}/\langle p'=1 \rangle$ as well.

Similarly, equivalence relation $P \sim Q$, implied by $pp'P + 1 = Q$, generates relation $P = Q \in R_{p,qq'}$ and relation $P = (q')^i Q \in R_{p,q}$ for some $i \in \mathbb{N} \cup \{0\}$. Then $P = Q$ is a defining relation of $H_{p,q}/\langle q'=1 \rangle$ as well. Therefore relations $P \sim Q$, where $P, Q \approx 1$, that generate $\overline{R}_{p,qq'}$, also generate defining relations of $\overline{H}_{p,q}/\langle q'=1 \rangle$.

Thus, group $\overline{H}_{p,q}/\langle p'=1 \rangle$ is isomorphic to a quotient group of $\overline{H}_{pp',q}$ and $\overline{H}_{p,q}/\langle q'=1 \rangle$ is isomorphic to a quotient group of $\overline{H}_{p,qq'}$. $\square$

Property 6. Let $q' \in \text{Ker } \phi_{p,q} \cap \text{Ker } \phi_{p,qq'}$. Then $\overline{H}_{p,q}/\langle q'=1 \rangle \cong \overline{H}_{p,qq'}/\langle q'=1 \rangle$.

Proof. By Property 3, defining relations of groups $\overline{H}_{p,q}/\langle q'=1 \rangle$ and $\overline{H}_{p,qq'}/\langle q'=1 \rangle$ are generated by the same set of relations. Then $P = 1$ is a defining relation of $\overline{H}_{p,q}/\langle q'=1 \rangle$ if and only if it is a defining relation of $\overline{H}_{p,qq'}/\langle q'=1 \rangle$. Therefore, $\overline{H}_{p,q}/\langle q'=1 \rangle \cong \overline{H}_{p,qq'}/\langle q'=1 \rangle$, since $q = 1, q' = 1, qq' = 1$ are defining relations of both groups $\square$

3 Equivalence classes of division sequences

Let us get back to division sequences $\mathcal{C}_{p,q}^M$ over domain $M \in \{\mathbb{Z}_{\{\neq 0\}}, \mathbb{Z}_{\{>0\}}\}$. Free group presentations $H_{p,q}$ are related to equivalence classes defined by $\mathcal{C}_{p,q}^M$ as follows. Relation (2) restricts equivalence relation $\sim$ to positive integer numbers co-prime to q , and relation (3) ensures that for every $P \in M$ such that $P \sim 1$ we have $P = 1$ in $H_{p,q}$ (the inverse implication may not be true). Therefore, elements of $H_{p,q}$ that are not identity contain positive integer numbers that are not equivalent to 1 in the sense of $\sim$.

Corollary 4. Let $c, d \in M$. If c and d lie in different elements of $H_{p,q}$, then $c \not\sim d$. Therefore, the number of different equivalence classes of $\mathcal{C}_{p,q}^M$ is at least the order of $H_{p,q}$. $\square$

The structure of groups $\text{Ker } \phi_{p,q}$ is somewhat an indication of the structure of integers equivalent to 1 in $\mathcal{C}_{p,q}^M$.

Corollary 5. Members of equivalence classes of $\sim$ that lie in identity element of $H_{p,q}$ form a group under multiplication. $\square$

Corollary 6. If $\text{Ker } \phi_{p,q} = F$, then $\text{Ker } \phi_{p,q^n} = F$ for all $n \in \mathbb{N}$. $\square$

Let us give several examples of how the above properties can be applied to specific division sequences.

Example 1. If $\mathcal{C}_{p,q}^{\mathbb{Z}_{\{>0\}}}$ has just one equivalence class, then $H_{p,q}^{\mathbb{Q}_{\{\neq 0, \cdot\}}}$ has order ≤ 3 .

Proof. Let us assume that $H_{p,q}^{\mathbb{Q}_{\{\neq 0, \cdot\}}}$ has order ≥ 3 . Then there exist cosets $a\text{Ker } \phi_{p,q}$, $b\text{Ker } \phi_{p,q}$ and $c\text{Ker } \phi_{p,q}$ where $a, c, b \in \mathbb{Q}_{\{<0\}}$. In this case $ab, ac \in \mathbb{Q}_{\{>0\}}$ imply that $a\text{Ker } \phi_{p,q} \cdot b\text{Ker } \phi_{p,q} = ab\text{Ker } \phi_{p,q} = \text{Ker } \phi_{p,q}$ and similarly $a\text{Ker } \phi_{p,q} \cdot c\text{Ker } \phi_{p,q} = \text{Ker } \phi_{p,q}$ - a contradiction. $\square$

Example 2. If $\mathcal{C}_{p,q^n}^M$, $n \in \mathbb{N}$, has just one equivalence class, then $H_{p,q}^M \cong \{1\}$. $\square$

Example 3. If $\mathcal{C}_{pp',q}^M$ contains a single equivalence class, then $H_{p,q}^M \cong \{1\}$. $\square$

Example 4. Let $q' \sim 1$ in $\mathcal{C}_{p,q}^M$ and $\mathcal{C}_{p,qq'}^M$. If $\mathcal{C}_{p,q}^M$ contains single equivalence class, $H_{p,qq'}^M \cong \{1\}$. If $\mathcal{C}_{p,qq'}^M$ contains single equivalence class, $H_{p,q}^M \cong \{1\}$. $\square$

Example 5. $H_{7,16}^M$ and $H_{7,2}^M$ are isomorphic to quotient groups of each other as $8 \in \text{Ker } \phi_{7,2} \cap \text{Ker } \phi_{7,16}$.
 $H_{5,2}^M$ and $H_{5,12}^M$ are isomorphic to quotient groups of each other as $6 \in \text{Ker } \phi_{5,2} \cap \text{Ker } \phi_{5,12}$.
 $H_{3,2}^M$ and $H_{3,8}^M$ are isomorphic to quotient groups of each other as $4 \in \text{Ker } \phi_{3,2} \cap \text{Ker } \phi_{3,8}$. $\square$

Example 6. Let $q' \in \text{Ker } \phi_{p,q} \cap \text{Ker } \phi_{p,qq'}$. If $\mathcal{C}_{p,qq'}^M$ contains one equivalence class, then $a \approx 1$ for $a \in M$ in $\mathcal{C}_{p,q}^M$ only if $q'|a$.
If $\mathcal{C}_{p,q}^M$ contains one equivalence class, then $a \approx 1$ for $a \in M$ in $\mathcal{C}_{p,qq'}^M$ only if $q'|a$.

Proof. Let $\mathcal{C}_{p,qq'}^M$ contain a single equivalence class. Then $H_{p,qq'}^M \cong \{1\}$ and $\overline{H}_{p,qq'}^M \cong F$. We have

$$\overline{H}_{p,qq'}^M / \langle q'=1 \rangle \cong \overline{H}_{p,q'}^M / \langle q'=1 \rangle \cong F / \langle q'=1 \rangle$$

by Property 6. Therefore all members of M that are not equivalent to 1 are not co-prime to q' . Similar argument is applied for the case when $\mathcal{C}_{p,q}^M$ contains a single equivalence class. $\square$

References

- Magnus, Karrass, Solitar 1966. W. Magnus, A. Karrass, D. Solitar, Combinatorial group theory, J. Wiley, 1966.
Lyndon, Schupp 2001. Roger C. Lyndon and Paul E. Schupp, Combinatorial Group Theory, Springer, 2001.
Vinberg 1989. E. B. Vinberg, Linear Representations of Groups, Birkhäuser Verlag, Basel, 1989.